



Slugs

Quick Facts...

- Slugs can be one of the most damaging pests of vegetable and flower Gardens in Colorado.
- Slugs thrive under high moisture conditions. The most effective way to reduce slug problems is to reduce humidity in a Garden.
- Metaldehyde baits can help control slugs if they are applied correctly and at the proper time.
- Slugs can be attracted to fermenting materials, such as beer, and trapped.

Slugs are one of the most destructive and difficult pests to control. Seedlings of many vegetables and flowers are favored foods, and high populations of slugs can cause difficulties in establishing a crop. Slugs also feed on many fruits and vegetables prior to harvest. This preharvest feeding results in wounds that allow various fungi and bacteria to enter and spoil the crop. In addition, the slime trails produced by slugs can contaminate Garden produce.

Slug injury is sometimes difficult to diagnose because slugs hide during the day. Smaller leaves may be eaten entirely, while only the edges of larger leaves may be consumed. Thick leaves often are rasped from the leaf underside. The presence of slime trails often is the best indication of slug activity.

History and Habits

Slugs and snails are classified as gastropods. They are more similar to clams and mussels than to other common Garden pests such as Insects. Slugs feed on a variety of plant materials that they eat by rasping with specialized mouthparts. They feed mostly at night, although they occasionally are active on overcast or rainy days. Slugs seek shelter during the day in soil cracks and under debris. During hot, dry weather, they may be temporarily inactive.

Slugs lay small masses of eggs in soil cracks. The young slugs that hatch from these eggs travel through the soil and damage germinating seeds and root Crops. Under favorable conditions, slugs complete development in a few months. The most common species in Colorado often have two generations per year: one in spring, another in fall.

Cultural Controls

Slugs are mostly water and produce large amounts of protective mucous. Consequently, slugs are susceptible to drying. The level of a slug population depends on the moisture conditions in a Garden. Any practices that decrease moisture will reduce slug problems, although effects may not be seen immediately.

To eliminate potential shelter for slugs, remove surface debris in and around the Garden and avoid organic mulches (straw, grass clippings). Also, increase air movement around plants and reduce high moisture conditions with trellises and wider plant spacing.

Use drip irrigation, soaker lines or other techniques to limit water and decrease the humidity around plants. Overhead irrigation should be done early in the day to allow more time for leaves and soil to dry before the nightly activity of slugs.

Attractants and Traps

Slugs are attracted to chemicals produced by many fermenting materials. These materials can be used to make attractant traps. For example, pans of beer or sugar-water and yeast mixtures can effectively attract, trap and drown slugs. A single baiting can remain effective for several days, as long as sufficient liquid remains. However, because the range of such traps is only a few feet, they must be placed throughout the planting to significantly reduce slug populations. Alcohol is not an attractant to slugs.

Selective use of trap boards or moistened newspaper placed on the soil surface also can be used to concentrate slugs that seek shelter under them. Check these shelters every morning and kill any slugs found. When not regularly maintained, remove the traps.

Trap Crops also may be used. Certain plants seem to be favored by slugs, such as lettuce, calendula and beans. These may be planted to divert slugs from feeding on the main Crops. For further control, early every morning remove the slugs from the trap Crops or treat the trap Crops to kill the slugs.

Repellents and Barriers

Slugs often avoid travelling over acid, alkali or abrasive materials. Diatomaceous earth, wood ashes and similar materials placed around plants provide some protection. However, moisture reduces the effect of these treatments.

Salt also is toxic to slugs and direct applications of table salt to a slug can kill it. This technique has limited usefulness because excessive salt can affect plants. Certain metal ions also are highly repellent to slugs. Barriers of copper foil exclude slugs from greenhouse benches and raised bed plantings. Other copper-based materials, such as copper sulfate, repel slugs.

Molluscicides

Pesticides effective against slugs and snails are known as molluscicides. These often are different chemicals than those used to control Insects and other Garden pests. Slugs are not susceptible to poisoning by most insecticides.

Metaldehyde is the most commonly used molluscicide. It usually is sold in the form of slug baits. Liquid formulations (Deadline, Slug-it, etc.) also have become popular. Metaldehyde can be used around vegetable and fruit Crops as well as in flower Gardens or among nonfood Crops. Metaldehyde is attractive and poisonous to slugs. After feeding on it, the slugs become immobilized and start to produce large amounts of mucous. Death results from excessive loss of water.

Successful use of metaldehyde baits requires careful application and favorable weather conditions. Slugs often can recover from metaldehyde poisoning if high moisture conditions occur. Also, control is poor during cool periods because slugs are relatively inactive. Apply metaldehyde when the evening is warm and the forecast for the following day is hot and dry. Young slugs are more susceptible than older slugs.

The range of metaldehyde attraction is limited so spread it throughout the infested area. Use 10 or more pieces of bait (flakes, pellets, liquid drops) per square yard. The amount needed for effective control varies depending on the attractiveness of other food in the area.

Metaldehyde is rapidly inactivated by sunlight so spread it underneath leaves late in the day. Sprays of ammonia make excellent contact molluscicides. However, to be effective they must be applied directly to the exposed slugs. Because slugs normally feed at night, such treatments are limited to periods such as an overcast evening following a late afternoon rain shower. Household ammonia, diluted to a 5 to 10 percent concentration, is effective for this purpose.